



IBT BIOSERVICES

21 Firstfield RD * Suite 100
Gaithersburg, MD 20878
877-411-2041
Services@ibtbioservices.com

Mouse anti-lipoteichoic acid (LTA) mAb

Catalog #: 0270-001

Lot #: 1309002

Immunogen: *Staphylococcus epidermidis*, Hay strain (ATCC 55133).

Description: Mouse monoclonal antibody (mAb) reactive to *Staphylococcus lipoteichoic acid*.

Supplied: 200 µg is supplied in PBS at a concentration of 0.92 mg/mL.

Purification: Antibody is purified using immobilized protein A.

Clonality: Monoclonal of the IgG1 subtype.

Relevance: Lipoteichoic acid (LTA) is the major proinflammatory structure present within the cell wall layer of most gram-positive bacteria. It plays an important role in the initiation and progression of bacterial infection, inflammation, and septic shock. The antibody can be used for detection of LTA produced from *Staphylococcus* cells.

Recommended Dilutions:

Optimal use dilution should be determined for each individual application.

ELISA 0.1 – 1.0 µg/mL

Opsionization assay: use at 80-160 µg/ml (optimized for *Staph. epidermidis*, Hay strain).

Storage: 2-3 weeks +4°C, -20°C or colder for long term


Cross Reactivity: Based on ELISA, this mAb reacts with LTA from *S. epidermidis*, Hay strain (ATCC 55133) as well as clinical strains of *S. epidermidis* types I, II and III. The antibody also reacts with LTA from *S. aureus* types 5 and 8, *Streptococcus pyogenes*, *Streptococcus faecalis* and *Streptococcus mutans*. This mAb does not react with peptidoglycan from *S. aureus*, peptidoglycan rhamnose or pneumococcal polysaccharides. It also does not react with live *E. coli* or *H. influenzae* type B.

Intended for research use only, not for human, therapeutic, or diagnostic applications.

The buyer cannot sell or otherwise transfer this product for Commercial Purposes without written approval of Integrated BioTherapeutics, Inc.

Copyright 2014. Integrated BioTherapeutics, Inc. All rights reserved

ELISA Data:



ELISA plates were coated overnight at room temperature with 1 µg/ml of LTA-BSA (Fina Biosolutions). After washing, mouse monoclonal anti-LTA (0270-001) was added at the indicated dilutions and incubated for 30 minutes at room temperature. Bound anti-LTA mAb (0270-001) was detected using gamma-specific goat anti-mouse IgG-HRP (SouthernBiotech). The assay was developed using one-component TMB and TMB Stop Solution (SouthernBiotech).

Related Products:

S. aureus Lipoteichoic acid (LTA)
Catalog #: 1600-001

Anti- *S. aureus* Alpha-Hemolysin Polyclonal Antibody
Catalog # 04-0010

Anti-*S. aureus* SEB mAb for Western blotting
Catalog # 0220-013

Anti- *S. aureus* SEB mAb for ELISA, Western & Neutralization
Catalog # 0220-009

Anti- *S. aureus* Gamma Hemolysin B Polyclonal Antibody
Catalog # 0314-001

Anti- *S. aureus* Gamma Hemolysin C Polyclonal Antibody
Catalog # 0314-002

S. aureus Gamma Hemolysin B (Hlg B) recombinant, purified protein,
Catalog # 1403-001

S. aureus Gamma Hemolysin C (Hlg C) recombinant, purified protein,
Catalog # 1403-002

Anti- *S. aureus* PVL LukS Polyclonal Antibody
Catalog # 04-0009

Anti- *S. aureus* LukS-PV Monoclonal Antibody
Catalog # 0221-001

S. aureus LukS-PV recombinant, purified protein
Catalog # 0530-001

Anti- *S. aureus* PVL LukF Polyclonal Antibody
Catalog # 0312-002

S. aureus LukF-PV recombinant, purified protein
Catalog # 0530-002