

Recombinant Angola marburgvirus Glycoprotein minus the Transmembrane Region (MARV-Angola rGPΔTM)

Catalog #: 0506-015

Lot #: 1710008

Description: Mature, recombinant, His-tagged Angola marburgvirus Glycoprotein minus the transmembrane domain (MARV-Angola rGP Δ TM) is supplied as purified protein. MARV-Angola rGP Δ TM is produced in Sf9 insect cells using baculovirus for expression and is purified by FPLC.


Storage: 2-3 weeks at -20°C, -80°C long term

Size: 100 μg of protein supplied in PBS (supplemented with glycerol, arginine and glutamic acid) at a concentration of 0.325 mg/mL. The theoretical molecular weight of the protein is ~ 60 kDa including the His-tag, without glycosylation. Because of the highly glycosylated nature of this protein, migration in an SDS-PAGE gel is slowed resulting in broad, diffuse bands representing differing glycosylation forms.

Relevance: Recombinant glycoprotein provides a means for antibody development, control protein for testing, and a tool to enhance research.

Western Blot: Quality control testing demonstrates strong detection of GP null and GP2 under reduced conditions.

Related Products: IBT provides a wide array of anti-filovirus specific antibodies and other infectious disease reagents. Please see our website, www.ibtbioservices.com for more details.



(A) SDS-PAGE and stain demonstrating 5 μ g and 1 μ g (lane 1, 2 respectively) of MARV-Angola rGP Δ TM His-tag protein under denaturing and reducing conditions. MW denotes Novex Sharp prestained protein markers. (B) Western blot detection of MARV-Angola rGP Δ TM at 50 ng, 100 ng and 500 ng (lanes 1-3). MARV-Angola rGP Δ TM was detected using IBT's polyclonal antibody (catalog # 0303-007) at 0.5 μ g/mL and anti-rabbit IgG-HRP conjugate, followed by substrate.

ELISA Data

MARV-Angola rGPΔTM ng/well	OD 650 nm
800.000	3.748
400.000	3.519
200.000	3.370
100.000	3.462
50.000	3.381
25.000	3.174
12.500	2.866
6.250	2.536
3.125	1.933
1.563	1.278
0.781	0.790
0.391	0.372

Plate was coated with MARV-Angola rGP Δ TM starting at 800 ng/well, serially diluted in DPBS. Washed plate was detected using one dilution of a positive control serum, followed with anti-IgG HRP conjugate and TMB substrate. OD₆₅₀ is reported. Background of MARV-Angola rGP Δ TM coated plate without positive control serum was 0.073 OD₆₅₀.

Intended for research use only. Not for human, therapeutic, or diagnostic applications.

The buyer cannot sell or otherwise transfer this product for Commercial Purposes without written approval of Integrated BioTherapeutics, Inc.