

4 Research Court, Suite 300 Rockville, MD 20850 877-411-2041 Services@ibtbioservices.com

S. aureus recombinant LukE

Catalog #: 0530-004

Lot #: 1805008

Description: Purified, *Staphylococcus aureus* recombinant leukocidin-E (rLukE). The rLukE (tag free) is expressed in *E. coli* and purified by FPLC. The theoretical molecular weight of the protein is 34,819 Daltons.

Storage: 2-3 weeks at -20°C, -80°C long term

Size: 100 μ g of protein is supplied in PBS plus 5% glycerol at a concentration of 1.182 mg/mL. Protein demonstrates a molecular weight of approximately 35 kDa.

Relevance: This protein may be used in functional toxicity assays in combination with rLukD, or as a control protein in ELISA assays or Western blotting when detecting toxins produced by different strains of *S. aureus*.

Recommended Dilutions:

ELISA: Assay-dependent dilution.


WB: Assay-dependent dilution; internal QC demonstrates detection of 100 ng of rLukE protein using anti-LukS-PV polyclonal antibody (cat# 04-0009) by Western blot analysis.

Cytotoxicity assay: Cytotoxicity can be detected in human neutrophils when used in combination with rLukD.

For additional *S. aureus* products, please visit:


http://www.ibtbioservices.com/reagents/staphylococcus/

SDS-PAGE and Western Blot Detection



A) SDS-PAGE of rLukE: 1 μ g (lane 1) and 5 μ g (lane 2). (B) Western blot detection of rLukE at 100 ng, using IBT's anti-LukS-PV polyclonal antibody (Cat#04-0009) at 0.5 μ g/mL and an antirabbit IgG-HRP conjugate, followed by TMB membrane substrate.

Toxin Functionality



Human promyelocytic leukemia cell line HL60 was differentiated into neutrophils by treatment with DMSO. Neutrophils were incubated with serial dilutions of rLukE and rLukD at equimolar concentration for 3 hours at 37°C with 5% CO2 and 95% humidity. Cellular viability was determined by adding XTT and incubation for additional 16 hours. Cells were centrifuged and the OD determined in the supernatants at 470/690 nm. EC50 values were found to be 22.2 nM for the current lot 1805008 (red circles) and 17.5 nM for the previous lot 1503005 (blue circles).