

Human anti-LukF-PV mAb (051)

Catalog #: 0221-004

Lot #: 2204002

Immunogen: Panton-Valentine Leukocidin (PVL) F subunit

(LukF-PV)

Description: Human monoclonal antibody reactive to

 ${\it Staphylococcus \ aureus \ Luk F-PV}.$

Supplied: $100~\mu g$ is supplied in PBS at a concentration of 1.795

mg/mL.

Purification: Antibody is purified using immobilized protein G.

Storage: 2-3 weeks +4°C, ≤-20°C long term

Clonality: Monoclonal of the IgG1 subtype

Recommended Dilutions:

ELISA: Assay-dependent dilution. The antibody can be used for capture of LukF-PV in an indirect ELISA assay and Western blot detection.

WB: Suggested use dilution of 0.50 μg/mL


Neutralization: Assay-dependent dilution

Cross Reactivity: This antibody does not appear to cross react with S. aureus antigens Hla, LukS, SEA, or SEB, at the tested

concentrations.

For additional S. aureus products, please visit:

https://www.ibtbioservices.com/product-category/staphylococcus/



Western blot detection of Hla (lane 1), LukF (lane 2), LukS (lane3), SEA (lane 4) and SEB (lane 5), 200ng each. LukF-PV was detected with monoclonal antibody 051 at 0.50 μ g/mL and an antihuman IgG-HRP conjugate.

Capture ELISA Data:

ng/mL	OD 650 nm				
	Hla	LukF	LukS	SEA	SEB
100.000	0.111	3.594	0.109	0.094	0.100
10.000	0.114	3.642	0.081	0.072	0.071
1.000	0.108	3.759	0.086	0.085	0.080
0.100	0.091	3.615	0.085	0.078	0.0097
0.010	0.086	3.303	0.099	0.085	0.112
0.001	0.083	1.694	0.095	0.084	0.095
0.000	0.088	0.339	0.120	0.091	0.096
0.000	0.094	0.117	0.103	0.112	0.108

S. aureus antigens were diluted to 2 $\mu g/mL$ in PBS for plate coating. Anti-LukF-PV antibody 051 was serially diluted ten-fold from 100 $\mu g/mL$ and incubated on the coated plates. Washed plates were detected with anti-human IgG-HRP conjugate and TMB substrate. 0D $_{650}$ is reported above.